

BIDEN ADMINISTRATION PROPOSAL: REDUCTION OF NICOTINE CONTENT IN CIGARETTES

Natalie R. Beylin, BS; Anna Song, PhD

One of the preliminary proposed rules included in the Biden Administration's Spring 2022 Unified Agenda of Regulatory and Deregulatory Actions agenda is a "tobacco product standard that would establish a maximum nicotine level in cigarettes and certain finished tobacco products".¹

WHAT IS THE PURPOSE OF THIS RULE?

This initiative was proposed to reduce the amount of tobacco related deaths and illness.

- Over 480,000 people die in the United States per year from smoking-related disease, including 41,000 secondhand smoke deaths, and over 16 million Americans live with a smoking-related disease.²

HOW WILL IT WORK?

Nicotine is the addictive chemical in cigarettes and is the driving force behind the difficulties of quitting smoking.³

- More than half of adult cigarette smokers report having tried to quit in 2020.⁴
- Fewer than 10% of adult cigarette smokers succeed in quitting each year.⁴
- In 2021, 65.3% of youth who currently used tobacco reported wanting to quit.⁵

Thus, limiting the amount of nicotine in these products can help a) addicted smokers quit and b) to prevent youth from initiating use and becoming regular smokers.⁶

- Majority of adults favor reducing nicotine levels in cigarettes, including 8/10 smokers.⁷
- Simulation models show that reduced nicotine content in cigarettes would lead to significant reduction in tobacco-related deaths.⁸
 - o Estimated prevention of 2.8 million tobacco-related deaths by 2060.⁸

WOULD PEOPLE SMOKE MORE TO COMPENSATE FOR THE LOWER NICOTINE LEVELS?

- Results from clinical trials testing low-level reductions in nicotine to 0.4mg or less nicotine/g tobacco show participants smoking less cigarettes per day with less withdrawals and more motivation to quit.⁹⁻¹³
 - o Only moderate-level nicotine reductions, such as 5.2mg per gram of tobacco, have been reported to lead to compensatory smoking.^{11,12,14}
 - o Suggests nicotine reductions should be low-level; not moderate.



WHAT WILL BE THE MAXIMUM NICOTINE LEVEL IN CIGARETTES IF THIS PROPOSAL IS ENACTED?

- Although still undisclosed, FDA's Center for Tobacco Products has previously suggested a "sweet spot" range of 0.2- 0.7 mg nicotine per gram of tobacco cigarette to reduce amount of cigarettes smoked, addiction likelihood, dependence levels, and increased cessation rates without increased compensatory smoking, cravings, or withdrawals.¹⁵
- It is important that the maximum level set be not too high for several reasons:
 - Although varying amounts of reduced nicotine levels (2.4, 1.3, or 0.4mg) can contribute to decreased smoking frequency, quit attempts were much more likely to occur in individuals using the 0.4mg nicotine cigarettes.⁹
 - Some smokers still preferred the cigarettes with slightly higher levels of nicotine (2.4mg) per tobacco gram than the lowest level (0.4mg).¹⁶⁻¹⁸
 - Compensatory smoking only occurred in moderate level reductions.^{11,12,14}
- Research suggests a 0.4mg per gram of tobacco level would be most efficient and effective.¹⁹

WHAT ARE THE NEXT STEPS?

- FDA to issue the proposed rule by May 2023.²⁰

REFERENCES

1. Tobacco Product Standard for Nicotine Level of Certain Tobacco Products. Accessed June 25, 2022. <https://www.reginfo.gov/public/do/eAgendaViewRule?pubId=202204&RIN=0910-A176>
2. United States Surgeon General. The Health Consequences of Smoking – 50 Years of progress: A Report of the Surgeon General: (510072014-001). Published online 2014. doi:10.1037/e510072014-001
3. Abuse NI on D. Is nicotine addictive? National Institute on Drug Abuse. Published -. Accessed June 25, 2022. <https://nida.nih.gov/publications/research-reports/tobacco-nicotine-e-cigarettes/nicotine-addictive>
4. 4. USPHSO of the SG & National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. Smoking Cessation: A Report of the Surgeon General. Published online 2020:700.
5. Gentzke AS, Wang TW, Cornelius M, et al. Tobacco Product Use and Associated Factors Among Middle and High School Students – National Youth Tobacco Survey, United States, 2021. *MMWR Surveill Summ.* 2022;71(5):1-29. doi:10.15585/mmwr.ss7105a1
6. Commissioner O of the. FDA Announces Plans for Proposed Rule to Reduce Addictiveness of Cigarettes and Other Combusted Tobacco Products. FDA. Published June 24, 2022. Accessed June 25, 2022. <https://www.fda.gov/news-events/press-announcements/fda-announces-plans-proposed-rule-reduce-addictiveness-cigarettes-and-other-combusted-tobacco>
7. Ali FRM, Al-Shawaf M, Wang TW, King BA. U.S. Adults' Attitudes Toward Lowering Nicotine Levels in Cigarettes. *Am J Prev Med.* 2019;57(3):403-407. doi:10.1016/j.amepre.2019.04.016
8. Apelberg BJ, Feirman SP, Salazar E, et al. Potential Public Health Effects of Reducing Nicotine Levels in Cigarettes in the United States. *N Engl J Med.* 2018;378(18):1725-1733. doi:10.1056/NEJMs1714617
9. Randomized Trial of Reduced-Nicotine Standards for Cigarettes | NEJM. Accessed June 25, 2022. <https://www.nejm.org/doi/full/10.1056/nejmsa1502403>



REFERENCES

10. Nicotine Reduction in Cigarettes: Literature Review and Gap Analysis | Nicotine & Tobacco Research | Oxford Academic. Accessed June 25, 2022. https://academic.oup.com/ntr/article/21/Supplement_1/S133/5684964
11. Hatsukami DK, Kotlyar M, Hertsgaard LA, et al. Reduced nicotine content cigarettes: effects on toxicant exposure, dependence and cessation. *Addict Abingdon Engl.* 2010;105(2):343-355. doi:10.1111/j.1360-0443.2009.02780.x
12. Hatsukami DK, Luo X, Jensen JA, et al. Effect of Immediate vs Gradual Reduction in Nicotine Content of Cigarettes on Biomarkers of Smoke Exposure: A Randomized Clinical Trial. *JAMA.* 2018;320(9):880-891. doi:10.1001/jama.2018.11473
13. Smoking in the absence of nicotine: behavioral, subjective and physiological effects over 11 days - Donny - 2007 - *Addiction* - Wiley Online Library. Accessed June 25, 2022. <https://onlinelibrary.wiley.com/doi/full/10.1111/j.1360-0443.2006.01670.x>
14. Mercincavage M, Souprontchouk V, Tang KZ, et al. A Randomized Controlled Trial of Progressively Reduced Nicotine Content Cigarettes on Smoking Behaviors, Biomarkers of Exposure, and Subjective Ratings. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol.* 2016;25(7):1125-1133. doi:10.1158/1055-9965.EPI-15-1088
15. The Science to Inform a Tobacco Product Standard for the Level of Nicotine in Combusted Cigarettes - 10/11/2018 - 10/11/2018. FDA. Published May 22, 2019. Accessed June 25, 2022. <https://www.fda.gov/science-research/about-science-research-fda/science-inform-tobacco-product-standard-level-nicotine-combusted-cigarettes-10112018-10112018>
16. Higgins ST, Heil SH, Sigmon SC, et al. Addiction Potential of Cigarettes With Reduced Nicotine Content in Populations With Psychiatric Disorders and Other Vulnerabilities to Tobacco Addiction. *JAMA Psychiatry.* 2017;74(10):1056-1064. doi:10.1001/jamapsychiatry.2017.2355
17. Preliminary test of cigarette nicotine discrimination threshold in non-dependent versus dependent smokers - PubMed. Accessed June 25, 2022. <https://pubmed.ncbi.nlm.nih.gov/28380366/>
18. Perkins KA, Kunkle N, Karelitz JL. Threshold dose for behavioral discrimination of cigarette nicotine content in menthol vs. non-menthol smokers. *Psychopharmacology (Berl).* 2017;234(8):1255-1265. doi:10.1007/s00213-017-4563-3
19. White CM, Pickworth WB, Sved AF, Donny EC. Using Product Standards to Render the Most Harmful Tobacco Products Minimally Addictive: Maximum Nicotine Level, Non-Nicotine Constituents, and Scope. *Nicotine Tob Res.* 2019;21(Suppl 1):S13-S15. doi:10.1093/ntr/ntz121
20. Chappell B. The FDA knows nicotine is addictive. It wants to regulate it for the first time. NPR. <https://www.npr.org/2022/06/22/1106742729/fda-nicotine-level-regulate-cigarettes-tobacco>. Published June 22, 2022. Accessed June 25, 2022.

